

Cooks Pharmacy

Address: 101-520 University Ave Waterloo Phone: (519) 744-2200 Fax: (519) 744-2203

PATIENT INFORMATION			DH/	PHYSICIAN INFORMATION			
Name Date of Birth		Name			CPSO		
Addre	City	Phone	Addre	SS	City	Phone	
ANAL FISSURE							
	Hydrocortisone 150mg Lidocaine 200mg Ketoprofen 50mg Nitroglycerin 4mg Misoprostol 0.25mg Rectal Rocket – 6 suppositories Insert 1 suppository two-thirds of the way into the rectum (pointed end first) at bedtime for 3-6 consecutive nights						
	Diltiazem 2%, Hydrocortisone 2%, Lidocaine 3%, Nifedipine 0.2% Ointment – 50 grams Apply to the affected area three times a day and after each bowel movement						
Apply to the affected area times a day and after each bower movement							
HEMORRHOID							
	Hydrocortisone 150mg, Lidocaine 200mg Rectal Rocket – 6 suppositories Insert 1 suppository two-thirds of the way into the rectum (pointed end first) at bedtime for 3-6 consecutive nights						
	Zinc Oxide 10%, Hydrocortisone 2.5%, Phenylephrine 0.5%, Lidocaine 5%, Pramoxine 2% Ointment – 50 grams						
Apply to the affected area three times a day and after each bowel movement							
	ERECTILE DYSFUNCTION			HERPES LABIALIS			
	Sildenafil 50mg Rapid-Dissolve Tablets – 10 tablets			Acyclovir 5%, Deoxy-D-Glucose 0.2%, Ibuprofen 2%, Lidocaine 2%, Aloe Vera			
	Dissolve one tablet on the tongue at least 30 minutes before sexual activity as needed			0.2% Gel – 25 grams Apply to the affected area at least three times daily or as needed			
	Sildenafil 100mg Rapid-Dissolve Tablets – 10 tablets						
Ш	Dissolve one tablet on the tongue at least 30 minutes before sexual activity as needed			MIGRAINE			
URINARY TRACT INFECTION				Ketoprofen 12.5mg Riboflavin 200mg Caffeine Citrate 100mg Oral Capsules – 50 Capsules			
Phenazopyridine 200mg Capsules – 6 capsules				Take 2 capsules at the onset of migraine and every 4 hours as needed until pain subsides Sumatriptan 5% Ondansetron 3% Flurbiprofen 3% Ketamine 3% Gabapentin 2.5% Amitriptyline 2.5% Bupivacaine 1% Cream – 30 grams Apply to the back of the neck, at the hairline, at the onset of a migraine and			
Take 1 capsule three times a day after meals for 2 days							
NIPPLE SORENESS							
Miconazole 2%, Mupirocin 1%, Betamethasone 0.05% Ointment – 50 grams				as needed until pain subsides. May also apply to various trigger points on			
	Apply sparingly after each feeding. Do not wash or wipe off			the face.			
NAIL FUNGUS							
Ciclopirox 8%, Itraconazole 2% Ketoconazole 2% Terbinafine 2%, Ibuprofen 2%, Tea Tree Oil 5% Fungal Nail Lacquer – 15 millilitres							
Apply to adjacent skin and affected nails daily. Let dry. Remove with alcohol every 7 days. Repeat							
WARTS							
Salicylic Acid 15%, Cimetidine 10%, Lactic Acid 10%, Podophyllum Resin 10%, Deoxy-D-Glucose 0.25% Ointment – 30 grams Apply to the affected wart once daily then cover. Continue every 1 or 2 days until wart is resolved							
Salicylic Acid 60%, Deoxy-D-Glucose 0.25% in Petrolatum – 30 grams							
Apply to the affected wart once daily then cover. Continue every 1 or 2 days until wart is resolved							
ARTHRITIS & JOINT PAIN				NEUROPATHIC PAIN			
	Diclofenac 10% Cream – 100 grams Apply to the affected area three times a day			Ketoprofen 7.5%, Amitriptyline 5%, Lidocaine 5% Cream – 100 grams Apply to the affected area three times a day			
	Ketoprofen 12%, Ketamine 10% Cream – 100 grams			Ketamine 10%, Gabapentin 5%, Lidocaine 5% Cream – 100 grams			
<u> </u>	Apply to the affected area three times a day Ketamine 10%, Diclofenac 5%, Ketoprofen 11%	. Baclofen 2% Cream – 100		Apply to the affected area three Ketamine 10%, Amitriptyline 5	,	5%. Lidocaine 5% Cream –	
	grams	,		100 grams		,	
1	Apply to the affected area three times a day			Apply to the affected area thre	e times a dav		

Physician Signature Date